

October 25, 2023

1. This problem is to explain why we use the dipole–electric field interaction in the spectroscopy of molecules. The Hamiltonian for an atom in an external electromagnetic field is

$$H = \frac{1}{2m} [\hat{\mathbf{p}} - q\mathbf{A}(t, \hat{\mathbf{r}})]^2 + V(\hat{\mathbf{r}}). \quad (1)$$

Express it in the coordinate representation as

$$H = -\frac{1}{2m} [\nabla - iq\mathbf{A}(t, \mathbf{r})]^2 + V(\mathbf{r}). \quad (2)$$

The dipole approximation means within the atomic size the vector potential approximately remains constant with respect to \mathbf{r} , that is, $\mathbf{A}(t, \mathbf{r}) \simeq \mathbf{A}(t, \mathbf{r}_0) \equiv \mathbf{A}(t)$. Show that in the local gauge transformation of wave function,

$$\psi(t, \mathbf{r}) \rightarrow \psi'(t, \mathbf{r}) \equiv e^{-iq\mathbf{A}(t) \cdot \mathbf{r}} \psi(t, \mathbf{r}), \quad (3)$$

the Hamiltonian becomes

$$H \rightarrow H' = -\frac{\nabla^2}{2m} + V(\mathbf{r}) - \boldsymbol{\mu} \cdot \mathbf{E}(t), \quad (4)$$

with the Schrödinger equation remaining

$$i\dot{\psi}' = H'\psi'. \quad (5)$$

Here $\boldsymbol{\mu} \equiv q\mathbf{r}$ and $\dot{\mathbf{A}} = -\mathbf{E}$.